African Pegmatite (Pty) Ltd 27 Van Der Bijl Street, PO Box 598, Meyerton 1960 Gauteng, South Africa Email: wenger@pegmatite.co.za reception@pegmatite.co.za Website: www.mineralmilling.com

Tel: +27 016 362 0600/1/2/3/4 Int: +27 16 362 0600 Fax: 016 362 1239

## MILLERS & PROCESSORS OF BASE MINERALS



Approved By:
Compiled By: D. Genis

Data Sheet

Doc No. : DSL 003 Rev No. : 01 Date: 08 January 2007

Page 1 of 1

## LIME SULPHUR LIQUID CONCENTRATE

DIRECTIONS FOR USE (use only as directed) – Spray to wet all parts of plants thoroughly		
COMPATIBILITY – LIME SULPHUR is not compatible with most fungicides and insecticides		
CROP/DESEASE	DOSAGE	REMARKS
	(Per 100 litre	
	water)	
BOYSENBERRIES AND		
Youngberries	5.00 Litre	Apply at budswel
Anthracnose	2.50 Litre	Apply at green tip stage
CITRUS		
Flatmite, Greymite, Rustmite,	1.25 Litre	Full cover spray. Allow 4 weeks safety period between sulphur and oil treatments.
Budmite	1.25 Litre	Full cover spray. If severe blossom malformation; apply directly after blossoming and again in
		February. For light to medium malformation apply before April. Allow 4 weeks safety period
		between sulphur and oil treatment.
APRICOTS, PLUMS & PEACHES	8.00 to	Full cover dormant spray. It will also control overwintering eggs of Bryobia Mite, red and grey
Permicious Scale	12.00 Litre	scale.
ORNAMENTALS		
Permicious Scale	12.00 Litre	Full cover formant spray, on plants without leaves.
APPLES AND PEARS		
Permicious Scale	5.00 Litre plus	Full cover dormant sprays. It will also control overwintering of eggs of Bryobia, mite, grey scale,
	5.00 Litre winter oil	pear budmite, mealy bug and woolly apple aphid.
APPLES AND PEARS	2.50 Litre	Early green tip stage
Scab	2.00 Litre	Advanced green tip stage
Powdery Mildew	1.50 Litre	Pre-pink (Cluster) stage
	1.25 Litre	Pink stage (apples), Closed cluster (pears)
	1.00 Litre	Petal fall stage
	0.75 Litre	Thereafter do not use on Sulphur shy cultivars
PEACHES	5.000 Litre to	Apply as a single spray at budswell. Will assist in the control of powdery mildew. Dosage depends
Leaf Curl	7.00 Litre	on the sensitivity of cultivar.
VINES		
Dead arm, Erinose mite (budmite) and	12.00 Litre	Dormant spray. Apply 3 weeks before budswell.
Anthracnose		DI IOATION

## APPLICATION

An aqueous solution fungi-, acari- insecticide for the control of fungal diseases and pesta in crops as indicated. Active ingredient: Polysulphide sulphur......320g/l

WARNINGS - Handle with care. Skin, eye and mucous membrane irritant. Keep out of reach of children, uninformed persons and animals. Store away from food and feedstuffs. As LIME SULPHUR is inclined to cause burning when applied on cool, cloudy days as well as very hot days. It should not be applied on cloudy days nor when temperatures are in the vicinity of 30°C. LIME SULPHUR should not be used after green tip stage on Golden delicious, Ohenimuri and Granny Smith apples and Josephine and Doyenne Du Comice pears. Where LIME SULPHUR is used on citrus do not mix any oilspray within 4 weeks after applications of LIME SULPHUR.

PRECUATIONS – Do not inhale spray mist and avoid contact with skin and eyes. Wear protective gloves and face shield when handling concentrate. Wash contaminated clothing daily. Do not eat, drink or smoke whilst applying or before washing hands and face. Avoid spray drift onto other crops, grazing, rivers, dams and areas not under treatment. Clean application equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams. Invert the empty container over the spray or mixing tank to allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the risings to the contents of the spray tank before destroying the container. Destroy empty container and do not re-use for any other purposes.

The information and recommendations in this product data sheet are based on data we believe to be reliable. They are offered in good faith, but do not imply any warranty or performance guarantee, as conditions and methods of use of our products are beyond our control. As such, African Pegmatite Company South African makes no express or implied warranties of any kind with respect to this product, including but not limited to any implied warranty of merchantability or fitness for a particular purpose.

We recommend that the user determine whether the products and the information given are appropriate, and that the suitability and performance of our products are appropriate by testing with its own equipment. Specifications are subject to change without notice.

The information and recommendations given in this product data sheet should not be understood as a recommendation for the use of our products in violation of any patent or as a license to use any patents of the African Pegmatite Company South Africa.

Tel: +27 16 362-0600/4 Fax: +27 16 362-1239 Website: www.mineralmilling.com









